

RECEIVED 29 MAR 2002

Form PTO 1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (REV 5-93)		ATTORNEY'S DOCKET NUMBER P51034
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED / ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.5) 10/089433
INTERNATIONAL APPLICATION NO. PCT/US00/26951	INTERNATIONAL FILING DATE 29 September 2000	PRIORITY DATE CLAIMED 1 October 1999
TITLE OF INVENTION COMPOUNDS AND METHODS		
APPLICANT(S) FOR DO/EO/US Joseph P. Marino, Jr., Scott K. Thompson and Daniel F. Veber		

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98; and Form PTO-1449.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☒ Please amend the specification by inserting before the first line the sentence: This is a 371 of International Application PCT/US00/26951, filed September 29, 2000, which claims benefit from the following Provisional Application: 60/157,286, filed October 1, 1999.
16. ☐ A substitute specification.
17. ☐ A change of power of attorney and/or address letter.
18. ☒ An Abstract on a separate sheet of paper.
19. ☐ Other items or information:

US APPLICATION NO. (if known see 37 CFR 1.50) 10/089433		INTERNATIONAL APPLICATION NO. PCT/US00/26951		ATTORNEYS DOCKET NO. P51034	
20. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATIONS PTO USE ONLY	
Basic National Fee (37 C.F.R. 1.492(a)(1)-(5)):					
Search Report has been prepared by the EPO or JPO\$890.00					
International Preliminary Examination Fee paid to USPTO (37 CFR 1.492)\$710.00					
No International Preliminary Examination Fee paid to USPTO (37 CFR 1.492) but international search fee paid to USPTO (37 CFR 1.445(a)(2))\$740.00					
Neither International Preliminary Examination Fee (37 CFR 1.492) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$1,040.00					
International Preliminary Examination Fee paid to USPTO (37 CFR 1.492) and all claims satisfied provisions of PCT Article 33(2)-(4).....\$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$710.00	
Claims	Number Filed	Number Extra	Rate		
Total claims	21 - 20 =	1	1 x \$18.00	\$18.00	
Independent claims	5 - 3 =	2	2 x \$84.00	\$168.00	
Multiple dependent claims (if applicable)			+ \$280.00	\$0.00	
TOTAL OF ABOVE CALCULATIONS =				\$896.00	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$896.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)) +				\$	
TOTAL NATIONAL FEE =				\$896.00	
				Amount to be refunded	\$
				charged	\$

- a. ☐ A check in the amount of \$_____ to cover the above fees is enclosed.
- b. ☒ Please charge my Deposit Account No. 19-2570 in the amount of **\$896.00** to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-2570. A duplicate copy of this sheet is enclosed.
- d. ☒ General Authorization to charge any and all fees under 37 CFR 1.16 or 1.17, including petitions for extension of time relating to this application (37 CFR 1.136 (a)(3)).

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

GLAXOSMITHKLINE
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, PA 19406-0939
Phone (610) 270-5012
Facsimile (610) 270-5090

Kathryn L. Sieburth
SIGNATURE
Kathryn L. Sieburth
NAME
40,072
REGISTRATION NO.

10/089433

JCS RECEIVED 29 MAR 2002

PATENT

ATTORNEY'S DOCKET NUMBER P51034

TRANSMITTAL LETTER TO THE U.S. DESIGNATED OFFICE
(DO/US) - ENTRY INTO NATIONAL STAGE UNDER 35 USC 371

INTERNATIONAL APP. NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/US00/26951	29 September 2000	1 October 1999

TITLE OF INVENTION
COMPOUNDS AND METHODS

APPLICANT(S) FOR DO/US
Joseph P. Marino, Jr., Scott K. Thompson and Daniel F. Veber

Box PCT
Assistant Commissioner for Patents
Washington, D.C. 20231
ATTENTION: DO/US

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this Transmittal Letter, Form PTO 1390 and the papers indicated as being transmitted therewith, and Post Card are being deposited with the United States Postal Service on this date March 29, 2002 in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number EL421192718US addressed to the:

Assistant Commissioner for Patents, Washington, D.C. 20231.

Elsa Matos
(Typed or printed name of person mailing paper)

Elsa Matos
(Signature of person mailing paper)



20462

PATENT TRADEMARK OFFICE

10/089433

1010 Rec'd PCT/PTO 29 MAR 2002

"EXPRESS MAIL CERTIFICATE"

"EXPRESS MAIL" MAILING LABEL NUMBER EL421192718US

DATE OF DEPOSIT March 29, 2002

Attorney Docket No. P51034

INTERNATIONAL APP. NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/US00/26951	29 September 2000	1 October 1999

TITLE OF INVENTION
COMPOUNDS AND METHODS

APPLICANT(S) FOR DO/US

Joseph P. MARINO, Jr. Scott K. THOMPSON and Daniel F. VEBER

PRELIMINARY AMENDMENT

Preliminary to the examination of this application, Applicants respectfully request amendment of the above-identified application as follows:


In the Specification:

Kindly add the Abstract enclosed herewith on a separate sheet, at the end.

In the Claims:

Please amend Claim 16 as follows:

16. (Amended) A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof:



Formula (I)

wherein,

X is S or O;

R¹ is optionally substituted C₂-6alkyl, C₃-6alkenyl, C₃-6alkynyl, C₃-7cycloalkyl-C₀-6alkyl, optionally substituted Het-C₁alkyl or optionally substituted Ar-C₀-6alkyl, wherein Ar is an optionally substituted phenyl or naphthyl group;

R² is optionally substituted C₂-6alkyl, C₃-6alkenyl, C₃-6alkynyl, optionally substituted Ar-C₀-6alkyl, optionally substituted Het-C₀-6alkyl, C₃-7cycloalkyl-C₀-6alkyl, provided that when R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran,

Int'l App. No.: PCT/US00/26951
Int'l Filing Date: 29 September 2000

benzothiophene, benzisoxazole, benzothiozole or benzopyrazole, then the optional substituent is not $-(CH_2)_2NR^4R^5$; and

R^3 is H, optionally substituted C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, optionally substituted Ar- C_{0-6} alkyl, optionally substituted Het- C_{0-6} alkyl, or C_{3-7} cycloalkyl- C_{0-6} alkyl, C_{0-6} alkyl-C(O)X'AB, C_{0-6} alkyl-S(O) $_2$ X'AB, C_{0-6} alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, optionally substituted Ar- C_{0-6} alkyl, optionally substituted Het- C_{0-6} alkyl, C_{3-7} cycloalkyl- C_{0-6} alkyl, or A or B are independently absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

REMARKS


The above-identified application is being entered into the National Phase from PCT Application No. PCT/US00/26951.

The claims are 1-21, with claims 1, 6, 11, 16 and 21 being independent. Claim 16 has been amended. Support for this amendment may be found in the specification at page 7, lines 11-12 and at page 9, lines 21-22.

No new matter has been introduced.

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "**Version with markings to show changes made.**"

Respectfully submitted,




Kathryn L. Sieburth
Attorney for Applicants
Registration No. 40,072

SMITHKLINE BEECHAM CORPORATION
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, PA 19406-0939
Phone (610) 270-5012
Facsimile (610) 270-5090
n\kls\cases\P51034\prelim amd#2.doc

"Version with markings to show changes made."

16. (Amended) A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof:



Formula (I)

wherein,

X is S or O;

R¹ is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, [optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or] C₃₋₇cycloalkyl-C₀₋₆alkyl, optionally substituted Het-C₁alkyl or optionally substituted Ar-C₀₋₆alkyl, wherein Ar is an optionally substituted phenyl or naphthyl group;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, provided that when R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiazole or benzopyrazole, then the optional substituent is not -(CH₂)₂NR⁴R⁵; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

ABSTRACT OF THE DISCLOSURE

Compounds of this invention are non-peptide, reversible inhibitors of type 2 methionine aminopeptidase, useful in treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity.

COMPOUNDS AND METHODS

FIELD OF THE INVENTION

Compounds of this invention are non-peptide, reversible inhibitors of
5 type 2 methionine aminopeptidase, useful in treating conditions mediated by
angiogenesis, such as cancer, haemangioma, proliferative retinopathy,
rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular
neovascularization and obesity.

10 BACKGROUND OF THE INVENTION

In 1974, Folkman proposed that for tumors to grow beyond a critical
size and to spread to form metastases, they must recruit endothelial cells from
the surrounding stroma to form their own endogenous microcirculation in a
process termed angiogenesis (Folkman J. (1974) *Adv Cancer Res.* 19; 331).

15 The new blood vessels induced by tumor cells as their life-line of oxygen and
nutrients also provide exits for cancer cells to spread to other parts of the
body. Inhibition of this process has been shown to effectively stop the
proliferation and metastasis of solid tumors. A drug that specifically inhibits
this process is known as an angiogenesis inhibitor.

20 Having emerged as a promising new strategy for the treatment of
cancer, the anti-angiogenesis therapy ("indirect attack") has several advantages
over the "direct attack" strategies. All the "direct attack" approaches such as
using DNA damaging drugs, antimetabolites, attacking the RAS pathway,
restoring p53, activating death programs, using aggressive T-cells, injecting
25 monoclonal antibodies and inhibiting telomerase, etc., inevitably result in the
selection of resistant tumor cells. Targeting the endothelial compartment of
tumors as in the "indirect attack", however, should avoid the resistance
problem because endothelial cells do not exhibit the same degree of genomic
instability as tumor cells. Moreover, anti-angiogenic therapy generally has
30 low toxicity due to the fact that normal endothelial cells are relatively
quiescent in the body and exhibit an extremely long turnover. Finally since
the "indirect attack" and "direct attack" target different cell types, there is a
great potential for a more effective combination therapy.

More than 300 angiogenesis inhibitors have been discovered, of which
35 about 31 agents are currently being tested in human trials in treatment of
cancers (Thompson, et al., (1999) *J Pathol* 187, 503). TNP-470, a
semisynthetic derivative of fumagillin of *Aspergillus fuigatus*, is among the

most potent inhibitors of angiogenesis. It acts by directly inhibiting endothelial cell growth and migration *in vitro and in vivo* (Ingber et al. (1990) *Nature* 348, 555). Fumagillin and TNP-470, have been shown to inhibit type 2 methionine aminopeptidase (hereinafter MetAP2) by irreversibly modifying its active site. The biochemical activity of fumagillin analogs has been shown to correlate to their inhibitory effect on the proliferation of human umbilical vein endothelial cells (HUVEC). Although the mechanism of the selective action of fumagillin and related compounds on MetAP2-mediated endothelial cell cytostatic effect has not yet been established, possible roles of MetAP2 in cell proliferation have been suggested.

First, hMetAP-2-catalyzed cleavage of the initiator methionine of proteins could be essential for releasing many proteins that, after myristoylation, function as important signaling cellular factors involved in cell proliferation. Proteins known to be myristoylated include the src family tyrosine kinases, the small GTPase ARF, the HIV protein nef and the α subunit of heterotrimeric G proteins. A recently published study has shown that the myristoylation of nitric oxide synthase, a membrane protein involved in cell apoptosis, was blocked by fumagillin (Yoshida, et al. (1998) *Cancer Res.* 58(16), 3751). This is proposed to be an indirect outcome of inhibition of MetAP2-catalyzed release of the glycine-terminal myristoylation substrate. Alternatively, MetAP enzymes are known to be important to the stability of proteins *in vivo* according to the "N-end rule" which suggests increased stability of methionine-cleaved proteins relative to their N-terminal methionine precursors (Varshavsky, A (1996) *Proc. Natl. Acad. Sci. U.S.A.* 93, 12142). Inhibition of hMetAP2 could result in abnormal presence or absence of some cellular proteins critical to the cell cycle.

Methionine aminopeptidases (MetAP) are ubiquitously distributed in all living organisms. They catalyze the removal of the initiator methionine from newly translated polypeptides using divalent metal ions as cofactors. Two distantly related MetAP enzymes, type 1 and type 2, are found in eukaryotes, which at least in yeast, are both required for normal growth; whereas only one single MetAP is found in eubacteria (type 1) and archaeobacteria (type 2). The N-terminal extension region distinguishes the methionine aminopeptidases in eukaryotes from those in procaryotes. A 64-amino acid sequence insertion (from residues 381 to 444 in hMetAP2) in the catalytic C-terminal domain distinguishes the MetAP-2 family from the MetAP-1 family. Despite the difference in the gene structure, all MetAP

enzymes appear to share a highly conserved catalytic scaffold termed "pita-bread" fold (Bazan, et al. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91, 2473), which contains six strictly conserved residues implicated in the coordination of the metal cofactors.


5 Mammalian type 2 methionine aminopeptidase has been identified as a bifunctional protein implicated by its ability to catalyze the cleavage of N-terminal methionine from nascent polypeptides (Bradshaw, et al (1998) *Trends Biochem. Sci.* 23, 263) and to associate with eukaryotic initiation factor 2 α (eIF-2 α) to prevent its phosphorylation (Ray, et al. (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89, 539). Both the genes of human and rat MetAP2 were cloned and have shown 92% sequence identity (Wu, et al. (1993) *J Biol. Chem.* 268, 10796; Li, X. & Chang, Y.-H. (1996) *Biochem. & Biophys. Res. Comm.* 227, 152). The N-terminal extension in these enzymes is highly charged and consists of two basic polylysine blocks and one aspartic acid block, which has
10 been speculated to be involved in the binding of eIF-2 α (Gupta, et al. (1993) in *Translational Regulation of Gene Expression 2* (Ilan, J., Ed.), pp. 405-431, Plenum Press, New York).

The anti-angiogenic compounds, fumagillin and its analogs, have been shown to specifically block the exo-aminopeptidase activity of hMetAP2
20 without interfering with the formation of the hMetAP2 : eIF2 α complex (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099). Fumagillin and its analogs inactivate the enzymatic activity of hMetAP2 with a high specificity, which is underscored by the lack of effect of these compounds on the closely related type 1 methionine
25 aminopeptidase (MetAP1) both *in vitro* and *in vivo* in yeast (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099). The extremely high potency (IC₅₀ < 1 nM) of these inhibitors appears to be due to the irreversible modification of the active site residue, His231, of hMetAP2 (Liu, et al. (1998) *Science* 282, 1324). Disturbance of MetAP2
30 activity *in vivo* impairs the normal growth of yeast (Griffith, et al., (1997) *Chem. Biol.* 4, 461; Sin, et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 6099; In-house data) as well as *Drosophila* (Cutforth & Gaul (1999) *Mech. Dev.* 82, 23). Most significantly, there appears to be a clear correlation between the inhibition effect of fumagillin related compounds against the enzymatic
35 activity of hMetAP2 *in vitro* and the suppression effect of these compounds against tumor-induced angiogenesis *in vivo* (Griffith, et al., (1997) *Chem. Biol.* 4, 461).

Cancer is the second leading cause of death in the U.S., exceeded only by heart disease. Despite recent successes in therapy against some forms of neoplastic disease, other forms continue to be refractory to treatment. Thus, cancer remains a leading cause of death and morbidity in the United States and elsewhere (Bailar and Gornik (1997) *N Engl J Med* 336, 1569). Inhibition of hMetAP2 provides a promising mechanism for the development of novel anti-angiogenic agents in the treatment of cancers. It has now been discovered that compounds of formulae (I) and (IA) are effective inhibitors of hMetAP2, and thus would be useful in treating conditions mediated by hMetAP2.

SUMMARY OF THE INVENTION

In one aspect, the present invention is to a compound of formula (I), or a pharmaceutically active salt or solvate thereof, and its use in treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity:



Formula (I)

wherein:

X is S or O;


R¹ is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, provided that when R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiazole or benzopyrazole, then the optional substituent is not -(CH₂)₂NR⁴R⁵; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-

S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently
 5 absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

10 In a second aspect, the present invention is to a method of treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity by administering a compound of formula (IA), or a pharmaceutically acceptable
 15 salt or solvate thereof



Formula (IA)

wherein,

X is S or O;

20 R¹ is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl;
 25

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are
 30 independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent.

In another aspect, the present invention is to a method of inhibiting
 35 MetAP2 in the treatment of angiogenesis-mediated diseases, all in mammals,

preferably humans, comprising administering to such mammal in need thereof, a compound of formula (IA), or a pharmaceutically active salt or solvate thereof.

In yet another aspect, the present invention is to pharmaceutical compositions comprising a compound of formula (I) and a pharmaceutically acceptable carrier therefor. In particular, the pharmaceutical compositions of the present invention are used for treating MetAP2-mediated diseases.

DETAILED DESCRIPTION OF THE INVENTION

It has now been discovered that substituted 1,2,4-triazoles of formulae (I) and (IA) are inhibitors of MetAP2. It has also now been discovered that selective inhibition of MetAP2 enzyme mechanisms by treatment with the inhibitors of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, represents a novel therapeutic and preventative approach to the treatment of a variety of disease states; including, but not limited to, cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity.

The term "C₁₋₆alkyl" as used herein at all occurrences means a substituted and unsubstituted, straight or branched chain radical of 1 to 6 carbon atoms, unless the chain length is limited thereto, including, but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl, pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. Any C₁₋₆alkyl group may be optionally substituted independently by one or more of OR⁴, R⁴, NR⁴R⁵. C₀alkyl means that no alkyl group is present in the moiety. Thus, Ar-C₀alkyl is equivalent to Ar.

As used herein at all occurrences, substituents R⁴, R⁵, and R⁶ are independently defined as C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, Ar-C₀₋₆alkyl, Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl.

The term "C₃₋₇cycloalkyl" as used herein at all occurrences means substituted or unsubstituted cyclic radicals having 3 to 7 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl radicals.

The term "C₃₋₆alkenyl" as used herein at all occurrences means an alkyl group of 3 to 6 carbons wherein a carbon-carbon single bond is replaced by a carbon-carbon double bond. C₃₋₆alkenyl includes 1-propene, 2-propene, 1-butene, 2-butene, isobutene and the several isomeric pentenes and hexenes. Both cis and trans isomers are included within the scope of this invention.

Any C₃₋₆alkenyl group may be optionally substituted independently by one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶,

5 O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃ or halogen.

The term "C₃₋₆alkynyl" as used herein at all occurrences means an alkyl group of 3 to 6 carbons wherein one carbon-carbon single bond is replaced by a carbon-carbon triple bond. C₃₋₆ alkynyl includes 1-propyne, 2-propyne, 1-butyne, 2-butyne, 3-butyne and the simple isomers of pentyne and hexyne.

The terms "Ar" or "aryl" as used herein interchangeably at all occurrences mean phenyl and naphthyl, optionally substituted by one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶, O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃ or halogen; in addition, Ph may be optionally substituted with one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen; Het' is defined as for Het, and may be optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen; or two C₁₋₆alkyl or C₁₋₆alkoxy groups may be combined to form a 5-7 membered, saturated or unsaturated ring, fused onto the Ar ring.

Suitably, for compounds of formula (I), when Ar is substituted by Ph or Het', then Ph or Het' are substituted with one or more of C₂₋₆alkyl, C₁₋₆alkoxy, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃ or halogen.

The terms "Het" or "heterocyclic" as used herein interchangeably at all occurrences, mean a stable 5- to 7-membered monocyclic, a stable 7- to 10-membered bicyclic, or a stable 11- to 18-membered tricyclic heterocyclic ring, all of which are either saturated or unsaturated, and consist of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure.

It will be understood that Het may be optionally substituted with one or more of Ph-C₀₋₆alkyl, Het'-C₀₋₆ alkyl, C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, Ph-C₀₋₆alkoxy, Het'-C₀₋₆alkoxy, OH, NR⁴R⁵, Het'-S-C₀₋₆alkyl, (CH₂)₁₋₆OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, (CH₂)₀₋₆CO₂R⁶,
 5 O(CH₂)₁₋₆CO₂ R⁶, (CH₂)₁₋₆SO₂, CF₃, OCF₃, CN, or halogen; Ph may be optionally substituted with one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen; and two C₁₋₆alkyl or C₁₋₆alkoxy groups may be combined to form a 5-7 membered ring, saturated or unsaturated, fused onto the Het ring. Preferred optional
 10 substituents on Het are C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆mercaptyl, halogen, CF₃, OCF₃, CN, or NR⁴R⁵.

Het' is defined as for Het and may be optionally substituted by one or more of C₁₋₆alkyl, C₁₋₆alkoxy, OH, (CH₂)₁₋₆NR⁴R⁵, O(CH₂)₁₋₆NR⁴R⁵, CO₂R⁶, CF₃, or halogen.

15 Examples of such heterocycles include, but are not limited to piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, pyridinyl, pyrazinyl, oxazolidinyl, oxazolynyl, oxazolyl, isoxazolyl, morpholinyl, thiazolidinyl,
 20 thiazolynyl, thiazolyl, quinuclidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, benzoxazolyl, furyl, pyranyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzoxazolyl, benzofuranyl, benzothiophenyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl, as well
 25 as triazolyl, thiadiazolyl, oxadiazolyl, isoxazolyl, isothiazolyl, imidazolyl, pyridazinyl, pyrimidinyl and triazinyl which are available by routine chemical synthesis and are stable.

Compounds of this invention of formula (I), do not include compounds wherein R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiozole or benzopyrazole,
 30 and the optional substituent is -(CH₂)₂NR⁴R⁵. The following compounds of this invention are known: 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole. Fromm et al., *Justus Liebigs Ann. Chem.*, 437 1924, 113. A compound of formula (I) wherein R¹ is benzyl,
 35 R² is phenyl and R³ is hydrogen is known.

Suitably, when moieties R^1 , R^2 , or R^3 are either optionally substituted Ar-C₀₋₆alkyl or optionally substituted Het-C₀₋₆alkyl, the moiety may be attached to the triazole substituent through the aromatic ring or through the alkyl chain.

5 Further, it will be understood that when a moiety is "optionally substituted" the moiety may have one or more optional substituents, each optional substituent being independently selected.

The terms "hetero" or "heteroatom" as used herein interchangeably at all occurrences mean oxygen, nitrogen and sulfur.

10 The terms "halo" or "halogen" as used herein interchangeably at all occurrences mean F, Cl, Br, and I.

Here and throughout this application the term C₀ denotes the absence of the substituent group immediately following; for instance, in the moiety ArC₀₋₆alkyl, when C is 0, the substituent is Ar, e.g., phenyl. Conversely, 15 when the moiety ArC₀₋₆alkyl is identified as a specific aromatic group, e.g., phenyl, it is understood that C is 0.

Suitably X is sulfur or oxygen. Preferably X is sulfur.


Suitably, R^1 is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl. Preferably R^1 is optionally substituted 20 Ar-C₀₋₆alkyl or optionally substituted Het-C₀₋₆alkyl. More preferably R^1 is optionally substituted Ar-C₁alkyl or optionally substituted Het-C₁alkyl. Most preferably R^1 is optionally substituted benzyl, optionally substituted methylfuran or optionally substituted methylthiophene. Preferably, when R^1 25 is Het-C₁alkyl, the alkyl chain is directly attached to moiety X.

Suitably, R^2 is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl. Preferably, R^2 is optionally substituted Ar-C₀₋₆alkyl. More preferably R^2 is optionally substituted Ar-C₀alkyl. Most 30 preferably R^2 is optionally substituted Ar-C₀alkyl, wherein the optional substituent is ortho C₁₋₆alkyl, preferably branched C₁₋₆alkyl, most preferably isopropyl.

Suitably, R^3 is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are 35 independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl,

optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent. Preferably R³ is hydrogen or C₀₋₆alkyl-C(O)X'AB. More preferably R³ is hydrogen or C₀₋₆alkyl-C(O)X'AB, wherein X' is oxygen and A is methyl or hydrogen and B is absent.

A preferred compound of this invention is a compound of formula (IB):



Formula (IB).

Suitably, pharmaceutically acceptable salts of formula (I) include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate, or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, palmitate, salicylate, and stearate.

The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. The stereocenters may be (R), (S) or any combination of R and S configuration, for example, (R,R), (R,S), (S,S) or (S,R). All of these compounds are within the scope of the present invention.

All compounds of formula (IA) specifically named herein are considered to be part of the invention disclosed herein. Among the compounds of the invention of formula (IA) are the following compounds:

- 3-anilino-5-benzylthio-1,2,4-triazole;
- 3-anilino-5-methylthio-1,2,4-triazole;
- 3-anilino-5-(4-chloro-benzylthio)-1,2,4-triazole;
- 3-anilino-5-allylthio-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole;
- 3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole;
- 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-anilino-5-(propyl acetylthio)-1,2,4-triazole;

- 3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole;
3-anilino-5-(2-phenethylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole;
5 3-anilino-5-(1*H*-benzoimidazol-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole;
10 3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole;
3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole;
4-(5-anilino-2 *H*-[1,2,4]triazol-3-yl thio)-benzoic acid;
3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole;
15 3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
20 3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
25 3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
30 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
35 3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

- 3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 5 3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 10 3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 15 3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 20 3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 4-(5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
 25 methyl ester;
 4-(5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
 methyl ester;
 4-(5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
 methyl ester;
 30 4-(5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl
 ester;
 4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-
 benzoic acid methyl ester;
 4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
 35 methyl ester;
 4-(5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl
 ester;

- 4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 5 4-(5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 4-(5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
- 3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
- 10 3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
- 15 3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
- 20 3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
- 25 3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
- 30 3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
- 3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
- [5-(benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- 35 [5-(3-methoxybenzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- [5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- [5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;

- [5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
- 5 [5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-
- 10 amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 15 3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
- 20 3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
- 25 3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
- 30 3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
- 35 3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;

- 3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
5 3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
10 3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
15 3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
20 3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
25 3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
30 3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
35 3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

- 3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
 5 3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 10 3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 15 3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
 20 3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 25 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 30 3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
 3-methyl-3-anilino-5-benzylthio-1,2,4-triazole;
 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
 3-*n*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
 35 3-*n*-butyl-3-anilino-5-benzylthio-1,2,4-triazole;
 3-*i*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
 3-allyl-3-anilino-5-benzylthio-1,2,4-triazole; and

3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole.

Among the preferred compounds of formula (IA) of this invention are the following compounds:

- 5 3-anilino-5-benzylthio-1,2,4-triazole;
- 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 10 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
- 3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
- 15 3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
- 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
- 20 3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
- 3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
- 3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
- 3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
- 3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
- 25 5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
- 5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
- 3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
- 30 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
- 3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
- 3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
- 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
- 35 3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
- 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;

- 3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
5 3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic
acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
10 carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
15 [5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
20 carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
25 3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
30 carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-
carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
35 3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;

- 3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 5 5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
 3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
 5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
 10 3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 15 3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 20 4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
 4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
 4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
 25 methyl ester;
 3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
 3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
 30 [5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
 [5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
 [5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
 [5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
 3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 35 3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;

- 3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 5 3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 10 3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 15 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
 3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
 3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole.

20


Among the more preferred compounds of formula (IA) are the following compounds:

- 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
 25 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
 4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
 3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
 3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
 30 3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
 3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
 3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
 35 3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;

- 3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
 3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 5 3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
 3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
 3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole.
- 10 Among the most preferred compounds of formula (IA) are the following compounds:
 3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 15 3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
 20 5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
 3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
 25 3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
 3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole.

Methods of Preparation

- 30 Compounds of the formulae (I) and (IA) wherein X is S and R³ is H, were prepared by methods analogous to those described in Scheme 1.


Scheme 1

- 5 a) Thiourea, NaOH, H₂O/CH₃CN; b) EtI, Et₃N, DMF; c) H₂NNH₂, EtOH; d)
 10 R¹X (X = halogen), K₂CO₃, DMF; e) ClCH₂OCH₂CH₃, NaH, THF; f)
 R³CH₂Br, NaH, DMF; g) TFA.

An isothiocyanate (such as phenyl isothiocyanate) (1-Scheme 1) was treated with thiourea and sodium hydroxide in acetonitrile/water to provide 2-Scheme 1, which was treated with iodoethane and triethylamine in DMF to afford 3-Scheme 1. Treatment of 3-Scheme 1 with hydrazine in ethanol provided 4-Scheme 1, which was treated with an alkyl halide (such as benzyl bromide or 4-chlorobenzyl chloride) and potassium carbonate in DMF to give 5-Scheme 1. Triazole 5-Scheme 1 is protected as the methoxy methylethyl ether to afford 6-Scheme 1. Alkylation of 6-Scheme 1 with an alkyl halide (such as methyl iodide, ethyl iodide, *i*-isobutyl iodide, *n*-propyl iodide, butyl iodide, allyl bromide, benzyl bromide, and methyl bromoacetate) afforded the desired tertiary amine 7-Scheme 1. Deprotection of the MOM-ether 7-Scheme 1 with trifluoroacetic acid (TFA) provided the desired product 8-Scheme 1.

Compounds of the formulae (I) and (IA) wherein X is O may be prepared by methods analogous to those described in Scheme 2.

Scheme 2



a) Thiourea, EtI, EtOH; b) NH_2NH_2 , EtOH c) 1,1'-Carbonyldiimidazole, EtOH; d) R^1X (X = halogen), K_2CO_3 , DMF; e) $\text{ClCH}_2\text{OCH}_2\text{CH}_3$, NaH, THF; f) $\text{R}^3\text{CH}_2\text{Br}$, NaH, DMF; g) TFA.

A thiourea (such as phenylthiourea) (8-Scheme 2) may be treated with ethyl iodide and refluxed in EtOH, and the resulting S-ethyl thiourea is then heated in the presence of hydrazine to provide 9-Scheme 2. The hydrazine 9-Scheme 2 is treated with carbonyldiimidazole and heated to afford 10-Scheme 2. Treatment of 10-Scheme 2 with an alkyl halide (such as benzyl bromide or 4-chlorobenzyl chloride) and potassium carbonate in DMF gives 11-Scheme 2. Triazole 11-Scheme 2 is protected as the methoxy methylethyl ether to afford 12-Scheme 2. Alkylation of 12-Scheme 2 with an alkyl halide (such as methyl iodide, ethyl iodide, *i*-isobutyl iodide, *n*-propyl iodide, butyl iodide, allyl bromide, benzyl bromide, and methyl bromoacetate) affords the desired tertiary amine 13-Scheme 2. Deprotection of the MOM-ether 13-Scheme 2 with trifluoroacetic acid (TFA) provides the desired product 14-Scheme 2.

Formulation of Pharmaceutical Compositions

The pharmaceutically effective compounds of this invention (and the pharmaceutically acceptable salts thereof) are administered in conventional dosage forms prepared by combining a compound of this invention ("active ingredient") in an amount sufficient to treat cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic

neovascularization, psoriasis, ocular neovascularization or obesity ("MetAp2-mediated disease states") with standard pharmaceutical carriers or diluents according to conventional procedures well known in the art. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl monostearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25 mg to about 1000 mg. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

The active ingredient may also be administered topically to a mammal in need of treatment or prophylaxis of MetAP2-mediated disease states. The amount of active ingredient required for therapeutic effect on topical administration will, of course, vary with the compound chosen, the nature and severity of the disease state being treated and the mammal undergoing treatment, and is ultimately at the discretion of the physician. A suitable dose of an active ingredient is 1.5 mg to 500 mg for topical administration, the most preferred dosage being 1 mg to 100 mg, for example 5 to 25 mg administered two or three times daily.

By topical administration is meant non-systemic administration and includes the application of the active ingredient externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye and nose, and where the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration.

While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The active ingredient may comprise, for topical administration, from 0.001% to

10% w/w, e.g. from 1% to 2% by weight of the formulation although it may comprise as much as 10% w/w but preferably not in excess of 5% w/w and more preferably from 0.1% to 1% w/w of the formulation.

5 The topical formulations of the present invention, both for veterinary and for human medical use, comprise an active ingredient together with one or more acceptable carrier(s) therefor and optionally any other therapeutic ingredient(s). The carrier(s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

10 Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

15 Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous or alcoholic solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C for half an hour. Alternatively, the
20 solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol
25 and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application
30 to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

35 Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise

hydrocarbons such as hard; soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol. The formulation may

5 incorporate any suitable surface-active agent such as an anionic, cationic or non-ionic surfactant such as esters or polyoxyethylene derivatives thereof.

Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

10 The active ingredient may also be administered by inhalation. By "inhalation" is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The daily dosage amount of the active ingredient administered by inhalation is from about
15 0.1 mg to about 100 mg per day, preferably about 1 mg to about 10 mg per day.

In one aspect, this invention relates to a method of treating cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization or obesity, all in mammals, preferably humans, which comprises administering to such
20 mammal an effective amount of a MetAP2 inhibitor, in particular, a compound of this invention.

By the term "treating" is meant either prophylactic or therapeutic therapy. Such compound can be administered to such mammal in a conventional dosage form prepared by combining the compound of this
25 invention with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The
30 compound is administered to a mammal in need of treatment for cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization or obesity, in an amount sufficient to decrease symptoms associated with these disease states. The route of administration may be oral or parenteral.

35 The term parenteral as used herein includes intravenous, intramuscular, subcutaneous, intra-rectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are

generally preferred. The daily parenteral dosage regimen will preferably be from about 30 mg to about 300 mg per day of active ingredient. The daily oral dosage regimen will preferably be from about 100 mg to about 2000 mg per day of active ingredient.

5 It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of this invention will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular mammal being treated, and that such optimums can be determined by conventional techniques. It will
10 also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of the compound given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

15 EXAMPLES

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. In the Examples, proton NMR spectra were performed upon a Bruker 400 MHz NMR spectrometer, unless
20 otherwise indicated.

Example 1

Preparation of 3-anilino-5-benzylthio-1,2,4-triazole

a) 1-Phenyl-2,4-dithiobiuret

To a stirring solution of NaOH (0.52 g, 13.1 mmol) in 60 mL of 10%
25 H₂O:CH₃CN was added thiourea (1.0 g, 13.1 mmol). The resulting suspension was heated to 40 °C for 20 min. and then cooled to RT. To this mixture was added phenylisothiocyanate (1.5 ml, 13.1 mmol), and the clear yellow solution was stirred overnight. After stirring for 12 h, 1 N HCl was added until a white precipitate formed. The precipitate was filtered, washed with H₂O, and dried
30 under vacuum to produce the title compound as a light yellow powder (0.78 g, 30%). ¹H-NMR (400MHz, d₆-DMSO) δ 7.25 (t, 2H, J=7.3 Hz), 7.40 (t, 2H, J=7.9 Hz), 7.56 (d, 1H, J=7.9 Hz), 9.13-9.29 (broad singlet, 1H), and 10.26-10.79 (broad singlet, 2H).

b) 2-Ethyl-1-phenyl-2-isodithiobiuret

35 To a stirring solution of the compound of Example 1(a) (150 mg, 0.70 mmol) in 4 mL of DMF was added triethylamine (57 uL, 0.70 mmol). The resulting yellow/green solution was stirred for 10 min at RT. To this solution

was added ethyl iodide (100 μ L, 0.70 mmol), and the reaction mixture was stirred for 2 h at RT. The yellow solution was poured into 20 mL of H₂O and extracted four times with EtOAc. The organic extracts were dried over Na₂SO₄, filtered, concentrated, and the crude residue was subjected to column chromatography (silica gel; ethyl acetate/hexane) to afford the title compound as a white crystalline solid (108 mg, 64%). ¹H-NMR (400MHz, d₆-DMSO) δ 1.22 (t, 3H, J=7.2 Hz), 2.96 (quartet, 2H, J=7.2 Hz), 6.85 (d, 1H, J=7.6 Hz), 7.16 (t, 1H, J=7.2 Hz), 7.29-7.41 (m, 3H), 8.27 (broad singlet, 1H), 9.89 (broad singlet, 1H), and 10.99 (broad singlet, 1H).

c) 3-anilino-5-mercapto-1,2,4-triazole

To a stirring solution of the compound of Example 1(b) in 2 mL of EtOH was added 50 μ L of anhydrous hydrazine. The reaction mixture was heated at 80 °C for 1 h, concentrated to dryness, and then purified by preparative HPLC to yield the title compound as a white solid (30 mg, 37%). MS (ESI) 190.90 (M-H)⁺.

d) 3-anilino-5-benzylthio-1,2,4-triazole

To a stirring solution of the compound of Example 1(c) (23 mg, 0.12 mmol) in 1.2 mL of DMF was added K₂CO₃ (17 mg, 0.12 mmol), followed by benzyl bromide (20 mg, 0.12 mmol). The mixture was stirred overnight, filtered, and purified by preparative HPLC to afford the title compound as a white solid (30 mg, 70%). ¹H-NMR (400MHz, d₆-DMSO) δ 9.30 (broad singlet, 1H), 7.47 (d, 2H, J=8.1 Hz), 7.39 (d, 2H, J=7.3 Hz), 7.31 (t, 2H, J=7.3 Hz), 7.23 (quartet, 3H, J=7.3 Hz), 6.82 (t, 1H, J=7.3 Hz), and 4.3 (s, 2H).

Example 2

Preparation of 3-anilino-5-(4-chlorobenzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except substituting 4-chlorobenzyl bromide for benzyl bromide in step 1(d), the title compound was prepared as a white solid. ¹H-NMR (400MHz, d₆-DMSO) δ 9.32 (broad singlet, 1H), 7.46 (d, 2H, J=7.8 Hz), 7.41 (d, 2H, J=8.4 Hz), 7.36 (d, 2H, J=8.4 Hz), 7.22 (t, 2H, J=7.8 Hz), 6.82 (t, 1H, J=7.24 Hz), and 4.33 (s, 2H).

Example 3

Preparation of 3-anilino-5-methylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except methyl iodide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 413.2 (2M+H)⁺.

Example 4

Preparation of 3-anilino-5-allylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except allyl bromide
5 was substituted for benzyl bromide in step 1(d), the title compound was
prepared as a white solid. MS (ESI) 233.0 (M+H)⁺.

Example 5

Preparation of 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole

10 Following the procedure of Example 1(a)-1(d), except 1-bromo-3-
methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 261.2 (M+H)⁺.

Example 6

Preparation of 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 1-bromo-3-
methylbutane was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 263.2 (M+H)⁺.

Example 7

Preparation of 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole

20 Following the procedure of Example 1(a)-1(d), except 5-bromo-2-
methyl-2-pentene was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

Example 8

Preparation of 3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except (1-bromoethyl)
benzene was substituted for benzyl bromide in step 1(d), the title compound
30 was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 9

Preparation of 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole

35 Following the procedure of Example 1(a)-1(d), except
bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 10

Preparation of 3-anilino-5-(propyl acetylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except propyl bromoacetate was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 293.2 (M+H)⁺.

Example 11

Preparation of 3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-bromo-1,1-dimethoxy-propane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 295.2 (M+H)⁺.

Example 12

Preparation of 3-anilino-5-(2-phenethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except (2-bromoethyl)benzene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 13

Preparation of 3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 288.2 (M+H)⁺.

Example 14

Preparation of 3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-chloromethyl-5-phenyl-1,2,4-oxadiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 351.2 (M+H)⁺.

Example 15

Preparation of 3-anilino-5-(1H-benzimidazol-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-(chloromethyl)-benzimidazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2 (M+H)⁺.

Example 16

Preparation of 3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except 4-chloromethyl-2-(4-chlorophenyl)thiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 400.0 (M+H)⁺.

Example 17

Preparation of 3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

10 Following the procedure of Example 1(a)-1(d), except 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 304.2 (M+H)⁺.

Example 18

Preparation of 3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 284.2 (M+H)⁺.

20

Example 19

Preparation of 3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 284.0 (M+H)⁺.

Example 20

Preparation of 3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole

30 Following the procedure of Example 1(a)-1(d), except 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 21

Preparation of 3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole

35 Following the procedure of Example 1(a)-1(d), except 4-isopropylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 22Preparation of 3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole

- Following the procedure of Example 1(a)-1(d), except 8-bromomethylquinoline was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 334.2 (M+H)⁺.

Example 23Preparation of 3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole

- Following the procedure of Example 1(a)-1(d), except 4-acetamidobenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 340.2 (M+H)⁺.

Example 24

- Preparation of 4-(5-anilino-2 H-[1,2,4]triazol-3-yl thio)-benzoic acid

Following the procedure of Example 1(a)-1(d), except 4-(chloromethyl)benzoic acid was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 25Preparation of 3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 297.0 (M+H)⁺.

Example 26Preparation of 3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole

- Following the procedure of Example 1(a)-1(d), except 4-(trifluoromethyl)benzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 350.8 (M+H)⁺.

Example 27Preparation of 3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole

- Following the procedure of Example 1(a)-1(d), except 3,5-dimethylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

Example 28Preparation of 3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-cyanobenzyl
5 bromide was substituted for benzyl bromide in step 1(d), the title compound
was prepared as a white solid. MS (ESI) 308.2 (M+H)⁺.

Example 29Preparation of 3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

10 Following the procedure of Example 1(a)-1(d), except 3,4-
difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title
compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 30Preparation of 3-anilino-5-(furan-2-ylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. J.
Chem. Soc. Perkin Trans. 1 1997, 8, 1147) was substituted for benzyl bromide
in step 1(d), the title compound was prepared as a white solid. MS (ESI)
20 273.2 (M+H)⁺.

Example 31Preparation of 3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R.
C. Tetrahedron Lett. 1988, 29(1), 117) was substituted for benzyl bromide in
step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2
(M+H)⁺.

Example 32Preparation of 3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

30 Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-
3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R.
C. Tetrahedron Lett. 1988, 29(1), 117) was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2
(M+H)⁺.

Example 33

Preparation of 3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 34

Preparation of 3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.0 (M+H)⁺.

Example 35

Preparation of 5-(5-phenylamino-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M)⁺.

Example 36

Preparation of 3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 367.0 (M)⁺.

Example 37

Preparation of 5-(5-phenylamino-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 301.2 (M+H)⁺.

Example 38

Preparation of 3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-chloromethylthiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 39

Preparation of 3-anilino-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-chloromethylfuran (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 273.2 (M+H)⁺.

Example 40

Preparation of 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 297.0 (M+H)⁺.

Example 41

Preparation of 3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 42

Preparation of 3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.0 (M+H)⁺.

Example 43Preparation of 3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 44Preparation of 3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

Example 45Preparation of 3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.0 (M+H)⁺.

Example 46Preparation of 3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 47Preparation of 3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.2 (M+H)⁺.

Example 48Preparation of 3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 49Preparation of 3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 50Preparation of 3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 51Preparation of 3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 52Preparation of 3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* 1997, 8, 1147) was substituted for benzyl

bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 53

5 Preparation of 3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 54

15 Preparation of 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 55

25 Preparation of 3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 56

35 Preparation of 3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-

chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

5

Example 57Preparation of 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

10

Example 58Preparation of 3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 381.0 (M)⁺.

20

Example 59Preparation of 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

30

Example 60Preparation of 3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.*

35

1958, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 61

5 Preparation of 3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* 1993, 10, 1941) was substituted for benzyl bromide in
10 step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 62

15 Preparation of 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 63

20 Preparation of 3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the
25 title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 64

30 Preparation of 3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 65

35 Preparation of 3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d) except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-

(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 66

5 Preparation of 3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

Example 67

10 Preparation of 3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 68

20 Preparation of 3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.0 (M+H)⁺.

Example 69

30 Preparation of 3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.2 (M+H)⁺.

Example 70

35 Preparation of 3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-

difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 71

Preparation of 3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 72

Preparation of 3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 73

Preparation of 3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 74

Preparation of 3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 75Preparation of 3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 76Preparation of 3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 77Preparation of 3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. J. *Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 78Preparation of 3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 79Preparation of 5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

5 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 80Preparation of 3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 381.0 (M)⁺.

Example 81Preparation of 5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

25 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 82Preparation of 3-(2-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

30 Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 83

Preparation of 3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *o*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, *10*, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 84

Preparation of 3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 85

Preparation of 3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 322.7 (M)⁺.

Example 86

Preparation of 3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 323.2 (M+H)⁺.

Example 87

Preparation of 3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 88

Preparation of 3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 295.2 (M+H)⁺.

Example 89

Preparation of 3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 335.2 (M+H)⁺.

Example 90

Preparation of 3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 322.0 (M+H)⁺.

Example 91

Preparation of 3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.0 (M+H)⁺.

Example 92

Preparation of 3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 352.8 (M)⁺.

Example 93Preparation of 3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 94Preparation of 3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 338.2 (M+H)⁺.

Example 95Preparation of 3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-chlorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.0 (M+H)⁺.

Example 96Preparation of 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 313.2 (M+H)⁺.

Example 97Preparation of 3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 98Preparation of 3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.2 (M+H)⁺.

Example 99Preparation of 3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 100Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 101Preparation of 3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 102Preparation of 3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 318.2 (M+H)⁺.

Example 103

Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 104

Preparation of 3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 105

Preparation of 3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 106

Preparation of 3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl

bromide in step 1(d), the title compound was prepared as a white solid (2%).
MS (ESI) 334.2 (M+H)⁺.

Example 107

5 Preparation of 3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2
10 (M+H)⁺.

Example 108

Preparation of 3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared
20 as a white solid . MS (ESI) 353.2 (M+H)⁺.

Example 109

Preparation of 3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except *p*-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid . MS (ESI) 353.0 (M+H)⁺.

30

Example 110

Preparation of 4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 341.0 (M+H)⁺.

Example 111Preparation of 4-(5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

5 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.2 (M+H)⁺.

Example 112Preparation of 4-(5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 342.2 (M+H)⁺.

Example 113Preparation of 4-(5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

20 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 114Preparation of 4-(5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 115Preparation of 4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

5 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 346.0 (M+H)⁺.

10

Example 116Preparation of 4-(5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

15 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.0 (M+H)⁺.

20

Example 117Preparation of 4-(5-(3-methoxy-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

25 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 118

30 Preparation of 4-(5-(3,4-difluoro-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

35 Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 377.0 (M+H)⁺.

Example 119Preparation of 4-(5-(2-methoxy-benzylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 120Preparation of 4-(5-(2-methyl-thiazol-4-yl)methylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 362.0 (M+H)⁺.

Example 121Preparation of 4-(5-(pyridin-2-yl)methylthio)-1H-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester

Following the procedure of Example 1(a)-1(d), except *p*-methoxycarbonylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 342.2 (M+H)⁺.

Example 122Preparation of 3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 123Preparation of 3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxybenzyl chloride was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 124

Preparation of 3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.2 (M+H)⁺.

Example 125

Preparation of 3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 126

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.2 (M+H)⁺.

Example 127

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.2 (M+H)⁺.

Example 128

Preparation of 3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 348.2 (M+H)⁺.

Example 129

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 130

Preparation of 3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 379.0 (M+H)⁺.

Example 131

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.0 (M+H)⁺.

Example 132

Preparation of 3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 364.2 (M+H)⁺.

Example 133

Preparation of 3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.0 (M+H)⁺.

Example 134

Preparation of 3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 135

Preparation of 3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 136Preparation of 3-(2-phenyl-anilino)-5-(3-methoxybenzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
5 isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-
methoxyphenyl chloride was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 389.0 (M+H)⁺.

Example 137

10 Preparation of 3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and
bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

15

Example 138Preparation of 3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d) except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-
20 (chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 360.2 (M+H)⁺.

Example 139Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-
bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the
title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

30

Example 140Preparation of 3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl
isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-
fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the
35 compound was prepared as a white solid. MS (ESI) 376.8 (M)⁺.

Example 141Preparation of 3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 364.0 (M+H)⁺.

Example 142Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.0 (M+H)⁺.

Example 143Preparation of 3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 395.0 (M+H)⁺.

Example 144Preparation of 3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 389.2 (M+H)⁺.

Example 145Preparation of 3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step

1(d), the title compound was prepared as a white solid. MS (ESI) 380.0 (M+H)⁺.

Example 146

Preparation of 3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-phenyl-phenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

Example 147

Preparation of [5-(benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 284.2 (M+H)⁺.

Example 148

Preparation of [5-(3-methoxybenzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-methoxyphenyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 149

Preparation of [5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 290.2 (M+H)⁺.

Example 150

Preparation of [5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-

(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 285.2 (M+H)⁺.

Example 151

5 Preparation of [5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the
10 title compound was prepared as a white solid. MS (ESI) 262.0 (M+H)⁺.

Example 152

Preparation of [5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

15 Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 153

20 Preparation of [5-(5-methyl-isoxazol-3-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step
25 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 154

30 Preparation of [5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the
35 title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 155Preparation of [5-(3,4-difluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 320.2 (M+H)⁺.

Example 156Preparation of [5-(2-methoxy-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 157Preparation of [5-(pyridin-2-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 285.2 (M+H)⁺.

Example 158Preparation of [5-(2-methyl-thiazol-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-chloromethyl-2-methylthiazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 305.2 (M+H)⁺.

Example 159Preparation of [5-(thiophen-2-ylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine

Following the procedure of Example 1(a)-1(d), except 3-pyridyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 290.0 (M+H)⁺.

5

Example 160

Preparation of 3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

10

Example 161

Preparation of 3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

15

Example 162

Preparation of 3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

20

Example 163

Preparation of 3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

30

Example 164

Preparation of 3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-

35

bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 289.0 (M+H)⁺.

Example 165

5 Preparation of 3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2.(M+H)⁺.

10

Example 166

Preparation of 3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

15

Example 167

20 Preparation of 3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

25

Example 168

Preparation of 3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

30

Example 169

35 Preparation of 3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-

methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.0 (M+H)⁺.

5

Example 170

Preparation of 3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 316.0 (M+H)⁺.

15

Example 171

Preparation of 3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 312.2 (M+H)⁺.

20

Example 172

Preparation of 3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-ethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

25

Example 173

Preparation of 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 313.2 (M+H)⁺.

35

Example 174

Preparation of 3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 175

Preparation of 3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 176

Preparation of 3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.2 (M+H)⁺.

Example 177

Preparation of 3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 178

Preparation of 3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide

in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 179

5 Preparation of 3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 180

15 Preparation of 3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 181

20 Preparation of 3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 182

30 Preparation of 3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

35

Example 183Preparation of 3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.0 (M+H)⁺.

Example 184Preparation of 3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 316.0 (M+H)⁺.

Example 185Preparation of 3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 314.2 (M+H)⁺.

Example 186Preparation of 3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid (49%). MS (ESI) 343.0 (M+H)⁺.

Example 187Preparation of 3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 188

Preparation of 3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 189

Preparation of 3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.2 (M+H)⁺.

Example 190

Preparation of 3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 191Preparation of 3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.0 (M+H)⁺.

Example 192Preparation of 5-(5-(2-methoxyphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 375.0 (M+H)⁺.

Example 193Preparation of 3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 396.8 (M-H)⁺.

Example 194Preparation of 3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 195Preparation of 3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 196Preparation of 3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 197Preparation of 3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 198Preparation of 3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 199Preparation of 3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in

step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 200

5 Preparation of 3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide
10 in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.2 (M+H)⁺.

Example 201

15 Preparation of 3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI)
20 303.2 (M+H)⁺.

Example 202

Preparation of 3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 203

30 Preparation of 3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 339.2 (M+H)⁺.

Example 204Preparation of 3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 205Preparation of 3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.2 (M+H)⁺.

Example 206Preparation of 3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

Example 207Preparation of 3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylisothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 330.2 (M+H)⁺.

Example 208Preparation of 3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 326.2 (M+H)⁺.

Example 209Preparation of 3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 355.2 (M+H)⁺.

Example 210Preparation of 3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Wish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 211Preparation of 3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

Example 212

Preparation of 3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

10 Example 213

Preparation of 3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

20 Example 214

Preparation of 3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 365.2 (M+H)⁺.

Example 215

30 Preparation of 5-(5-(2-isopropylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 387.2 (M+H)⁺.

Example 216

Preparation of 5-(5-(2-isopropyl amino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, 6, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 217

Preparation of 3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 218

Preparation of 3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-isopropylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 219

Preparation of 3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a), the title compound was prepared as a white solid. MS (ESI) 297.2 (M+H)⁺.

Example 220

Preparation of 3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-

chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 221

Preparation of 3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 222

Preparation of 3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 223

Preparation of 3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 275.2 (M+H)⁺.

Example 224

Preparation of 3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 225

Preparation of 3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step

1(d), the title compound was prepared as a white solid. MS (ESI) 302.2 (M+H)⁺.

Example 226

Preparation of 3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

Example 227

Preparation of 3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 228

Preparation of 3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 229

Preparation of 3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 331.2 (M+H)⁺.

Example 230

Preparation of 3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 231

Preparation of 3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

10

Example 232

Preparation of 3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

 Following the procedure of Example 1(a)-1(d), except *m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the
15 title compound was prepared as a white solid. MS (ESI) 298.2 (M+H)⁺.

Example 233

Preparation of 3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

20 Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Whish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS
25 (ESI) 287.2 (M+H)⁺.

Example 234

Preparation of 3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

30 Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS
35 (ESI) 317.2 (M+H)⁺.

Example 235

Preparation of 3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 236

Preparation of 3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, **1993**, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 237

Preparation of 3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 337.2 (M+H)⁺.

Example 238

Preparation of 5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 359.2 (M+H)⁺.

Example 239Preparation of 3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* **1947**, *60*, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 381.0 (M+H)⁺.

Example 240Preparation of 5-(5-(3-methylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, *6*, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 315.2 (M+H)⁺.

Example 241Preparation of 3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 303.2 (M+H)⁺.

Example 242Preparation of 3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 3-*m*-tolyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, *10*, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 287.2 (M+H)⁺.

Example 243Preparation of 3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 339.2 (M+H)⁺.

Example 244Preparation of 3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 345.2 (M+H)⁺.

Example 245Preparation of 3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 246Preparation of 3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 375.2 (M+H)⁺.

Example 247Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 317.2 (M+H)⁺.

Example 248Preparation of 3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

5

Example 249

Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 353.2 (M+H)⁺.

10

Example 250

Preparation of 3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

15

Example 251

Preparation of 3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

20

Example 252

Preparation of 3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 383.2 (M+H)⁺.

30

35

Example 253Preparation of 3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 254Preparation of 3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 340.2 (M+H)⁺.

Example 255Preparation of 3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-*n*-butylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 369.2 (M+H)⁺.

Example 256Preparation of 3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 343.0 (M+H)⁺.

Example 257Preparation of 3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1 (d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

Example 258Preparation of 3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

5 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

10

Example 259Preparation of 3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

15 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 349.0 (M+H)⁺.

20

Example 260Preparation of 3-(2,4-dimethoxy-anilino)- (3,4-difluoro-benzylthio)-1,2,4-triazole

25 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 379.0 (M+H)⁺.

Example 261

30 Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

35 Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.2 (M+H)⁺.

Example 262Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

Example 263Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 264Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 377.0 (M+H)⁺.

Example 265Preparation of 3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 266Preparation of 3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 387.2 (M+H)⁺.

Example 267Preparation of 3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 348.0 (M+H)⁺.

Example 268Preparation of 3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 344.2 (M+H)⁺.

Example 269Preparation of 3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,4-dimethoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 373.2 (M+H)⁺.

Example 270Preparation of 3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 327.2 (M+H)⁺.

Example 271Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethylthiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 272Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 273Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 333.2 (M+H)⁺.

Example 274Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 363.0 (M+H)⁺.

Example 275

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 305.2 (M+H)⁺.

Example 276

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 277

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 278

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 361.0 (M+H)⁺.

Example 279

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 280

Preparation 3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-methylenedioxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 371.0 (M+H)⁺.

Example 281

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-(chloromethyl)-5-methylisoxazole was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 332.2 (M+H)⁺.

Example 282

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-(chloromethyl)pyridine was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 328.2 (M+H)⁺.

Example 283

Preparation of 3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2-methyl-4-methoxyphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methoxybenzyl chloride was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 357.2 (M+H)⁺.

Example 284

Preparation of 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) the title compound was prepared as a white solid. MS (ESI) 311.4 (M+H)⁺.

Example 285

Preparation of 3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 4-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

Example 286

Preparation of 3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and bromomethylcyclohexane was substituted for benzyl bromide in

step 1(d), the title compound was prepared as a white solid . MS (ESI) 317.2 (M+H)⁺.

Example 287

5 Preparation of 3-(2,6-dimethyl-anilino)-(3,4-difluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3,4-difluorobenzyl bromide was substituted for benzyl bromide
10 in step 1(d), the title compound was prepared as a white solid. MS (ESI) 347.0 (M+H)⁺.

Example 288

15 Preparation of 3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 1-bromo-3-methylbut-2-ene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI)
20 289.0 (M+H)⁺.

Example 289

Preparation of 3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in
25 step 1(a) and 2-fluorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 329.2 (M+H)⁺.

30 Example 290

Preparation of 3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-methylbenzyl bromide was substituted for benzyl bromide in
35 step 1(d), the title compound was prepared as a white solid. MS (ESI) 325.2 (M+H)⁺.

Example 291

Preparation of 3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 2,6-dimethylphenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chlorobenzyl bromide was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 345.0 (M+H)⁺.

Example 292

Preparation of 3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-furan (Berry, J. M.; Watson, C. Y.; Wish, W. J. D.; Threadgill, M. D. *J. Chem. Soc. Perkin Trans. 1* **1997**, 8, 1147) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 293

Preparation of 3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-methyl-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.0 (M+H)⁺.

Example 294

Preparation of 3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-3-chloro-thiophene (Chauhan, P. M. S.; Jenkins, G.; Walker, S. M.; Storr, R. C. *Tetrahedron Lett.* **1988**, 29(1), 117) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 295

Preparation of 3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-methyl-thiophene (Moradpour, A. *J. Chem. Soc. Perkin Trans. 1*, 1993, 1, 7) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 321.0 (M+H)⁺.

Example 296

Preparation of 3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-chloro-thiophene was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 341.2 (M+H)⁺.

Example 297

Preparation of 5-(5-(4-fluorophenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carboxylic acid ethyl ester was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 363.2 (M+H)⁺.

Example 298

Preparation of 3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 2-chloromethyl-5-bromo-thiophene (Clapp, R. C.; Clark, J. H.; Vaughan, J. R.; English, J. P.; Anderson, G. W. *J. Am. Chem. Soc.* 1947, 69, 1549) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 385.0 (M)⁺.

Example 299

Preparation of 5-(5-(4-fluorophenylamino)-4H-[1,2,4]triazol-3-yl)sulfanylmethyl)-furan-2-carbaldehyde

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 5-chloromethyl-furan-2-carbaldehyde (Sanda, K.; Rigal, L.; Delmas, M.; Gaset, A. *Synthesis* **1992**, 6, 541) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 319.0 (M+H)⁺.

Example 300

Preparation of 3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-thiophene (Lamy, J.; Lavit, D.; Buu-Hoi, N. P. *J. Chem. Soc.* **1958**, 4202) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 307.2 (M+H)⁺.

Example 301

Preparation of 3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole

Following the procedure of Example 1(a)-1(d), except 4-fluorophenyl isothiocyanate was substituted for phenylisothiocyanate in step 1(a) and 3-chloromethyl-furan (Arena, G.; Cali, R.; Maccarone, E.; Passerini, A. *J. Chem. Soc. Perkin Trans. 2* **1993**, 10, 1941) was substituted for benzyl bromide in step 1(d), the title compound was prepared as a white solid. MS (ESI) 291.2 (M+H)⁺.

Example 302

Preparation of 3-methyl-3-anilino-5-benzylthio-1,2,4-triazole

a) 3-anilino-5-benzylthio-1 or/2-methyl ethyl ether-1,2,4-triazole

To a stirring solution of 3-anilino-5-benzylthio-1,2,4-triazole (0.68 g, 2.41 mmol) in 8 mL DMF was added NaH (0.125 g, 3.13 mmol). To this mixture was added chloromethyl ethyl ether (0.251 g, 2.65 mmol), and the solution was stirred overnight. The reaction mixture was poured into 50 ml H₂O and extracted three times with EtOAc. The EtOAc extracts were dried over Na₂SO₄, filtered, and concentrated down. The crude mixture was subjected to column chromatography (silica gel, EtOAc/hexane) to provide the title compounds as a mixture of regioisomers as a light yellow oil (0.58 g,

71%). ¹H-NMR (400MHz, d6-DMSO) compound 1: δ9.33 (broad singlet, 1H), 7.51 (d, 2H, J=8.3 Hz), 7.42-7.22 (m, 8H), 5.23 (s, 2H), 4.47 (s, 2H), 3.43 (q, 2H, J=7.2 Hz), 1.04 (t, 3H, J=7.0 Hz). Compound 2: δ9.20 (broad singlet, 1H), 7.63 (d, 2H, J=7.6 Hz), 7.42-6.93 (m, 8H), 5.44 (s, 2H), 4.30 (s, 2H), 3.51 (q, 2H, J=7.1 Hz), 1.07 (t, 3H, J=7.0). MS (ESI) 341 (M+H)⁺.

b) 3-methyl-3-anilino-5-benzylthio-1,2,4-triazole

To a stirring solution of 3-anilino-5-benzylthio-1 or/2-methyl ethyl ether-1,2,4-triazole (50 mg, 0.15 mmol) in 1 ml THF was added NaH (11.8 mg, 0.30 mmol), and to this solution was added CH₃I (0.036 ml, 0.57 mmol). The reaction mixture was stirred overnight. THF was removed and 0.5 ml TFA was added to the residue and stirred overnight. TFA was removed under vacuum and the mixture was purified by preparative HPLC to afford the title compound as a clear oil (28 mg, 53%). ¹H-NMR (400MHz, d6-DMSO) δ3.3-7.25 (m, 10H), 4.27 (s, 2H), 3.40 (s, 3H). MS (ESI) 297 (M+H)⁺.

Example 303

Preparation of 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except iodoethane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.26 (m, 10H), 4.26 (s, 2H), 3.86 (m, 2H), 1.20 (m, 3H). MS (ESI) 311 (M+H)⁺.

Example 304

Preparation of 3-n-propyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodopropane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (35%). ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.26 (m, 10H), 4.25 (s, 2H), 3.76 (t, 2H, J=6.5 Hz), 3.31 (t, 2H, J=1.4 Hz), 1.63 (m, 2H), 0.93 (t, 3H, J=7.4 Hz). MS (ESI) 325 (M+H)⁺.

Example 305

Preparation of 3-n-butyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodobutane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (31%). ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.22 (m, 10H), 4.26 (s, 2H), 3.80 (t, 2H, J=7.5 Hz), 3.31 (t, 2H, J=1.4 Hz), 1.59 (m, 2H), 1.36 (m, 2H), 0.92 (t, 3H, J=7.3 Hz). MS (ESI) 338 (M+H)⁺.

Example 306

Preparation of 3-isopropyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 1-iodo-2-methyl propane was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.42-7.22 (m, 10H), 4.25 (s, 2H), 3.66 (d, 2H, J=7.6 Hz), 1.92 (m, 1H), 0.93 (d, 6H, J=6.7 Hz). MS (ESI) 338 (M+H)⁺.

Example 307

Preparation of 3-allyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except allyl bromide was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (41%). ¹H-NMR (400MHz, d6-DMSO) δ7.37-7.28 (m, 10H), 5.96 (m, 1H), 5.18 (m, 2H), 4.45 (s, 2H), 4.26 (s, 2H). MS (ESI) 323 (M+H)⁺.

Example 308

Preparation of 3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except benzyl bromide was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (48%). ¹H-NMR (400MHz, d6-DMSO) δ7.28-7.23 (m, 15H), 5.09 (s, 2H), 4.26 (s, 2H). MS (ESI) 373 (M+H)⁺.

Example 309

Preparation of 3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid. ¹H-NMR (400MHz, d6-DMSO) δ7.37-7.22 (m, 10H), 4.59 (s, 2H), 4.26 (s, 2H), 3.74 (s, 3H). MS (ESI) 355 (M+H)⁺.

Example 310

Preparation of 3-methylacetate-3-(p-methyl)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(p-methyl)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-

anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a clear oil. ¹H-NMR (400MHz, d6-DMSO) δ7.38-7.09 (m, 9H), 4.56 (s, 2H), 4.27 (s, 2H), 3.75 (s, 3H), 2.37 (s, 3H). MS (ESI) 369 (M+H)⁺.

5

Example 311

Preparation of 3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a brown oil (44%). ¹H-NMR (400MHz, d6-DMSO) δ7.92-7.22 (m, 7H), 6.99 (d, 2H, J=8.9 Hz), 4.51 (s, 2H), 4.26 (s, 2H), 3.83 (s, 3H), 3.76 (s, 3H). MS (ESI) 385 (M+H)⁺.

15

Example 312

Preparation of 3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole

Following the procedure of Example 302(a)-(b) except 3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole was used in step 302(a) instead of 3-anilino-5-benzylthio-1,2,4-triazole and methyl bromoacetate was used in step 302(b) instead of iodomethane, the title compound was isolated as a white solid (43%). ¹H-NMR (400MHz, d6-DMSO) δ7.32-7.19 (m, 8H), 4.37 (s, 2H), 4.25 (s, 2H), 3.77 (s, 3H), 2.27 (s, 6H). MS (ESI) 383 (M+H)⁺.

25

Biological Data:

Direct Spectrophotometric Assays of hMetAP2:




The hMetAP2 activity can be measured by direct spectrophotometric assay methods using alternative substrates, L-methionine-*p*-nitroanilide (Met-pNA) and L-methionine-7-amido-4-methylcoumarin (Met-AMC). The formation of *p*-nitroaniline (pNA) or 7-amido-4-methylcoumarin (AMC) was continuously monitored by increasing absorbance or fluorescence at 405 nm and 460 nm, respectively, on a corresponding plate reader. All assays were carried out at 30°C. The fluorescence or spectrophotometric plate reader was calibrated using authentic pNA and AMC from Sigma, respectively. For a typical 96-well plate assay, the increase in the absorbance (at 405 nm for pNA) or the fluorescence

35

emission ($\lambda_{\text{ex}} = 360 \text{ nm}$, $\lambda_{\text{em}} = 460 \text{ nm}$, for AMC) of a 50 μL assay solution in each well was used to calculate the initial velocity of hMetAP2. Each 50 μL assay solution, contained 50 mM Hepes- Na^+ (pH 7.5), 100 mM NaCl, 10-100nM purified hMetAP2 enzyme, and varying amounts of Met-AMC (in 3% DMSO aqueous solution) or Met-pNA. Assays were initiated with the addition of substrate and the initial rates were corrected for the background rate determined in the absence of hMetAP2.

Coupled Spectrophotometric Assays of hMetAP2:

The methionine aminopeptidase activity of hMetAP2 can also be measured spectrophotometrically by monitoring the free L-amino acid formation. The release of N-terminal methionine from a tripeptide (Met-Ala-Ser, Sigma) or a tetrapeptide (Met-Gly-Met-Met, Sigma) substrate was assayed using the L-amino acid oxidase (AAO) / horse radish peroxidase (HRP) couple (eq. 1-3a,b). The formation of hydrogen peroxide (H_2O_2) was continuously monitored at 450nm (absorbance increase of o-Dianisidine (Sigma) upon oxidation, $\Delta\epsilon = 15,300 \text{ M}^{-1}\text{cm}^{-1}$)² and 30 °C in a 96- or 384-well plate reader by a method adapted from Tsunasawa, S. et al.(1997) (eq. 3a). Alternatively, formation of H_2O_2 was followed by monitoring the fluorescence emission increase at 587nm ($\Delta\epsilon = 54,000 \text{ M}^{-1}\text{cm}^{-1}$, $\lambda_{\text{ex}} = 563 \text{ nm}$, slit width for both excitation and emission was 1.25 mm) and 30 °C using Amplex Red (Molecular Probes, Inc) (Zhou, M. et al. (1997) *Anal. Biochem.* 253, 162) (eq. 3b). In a total volume of 50 μL , a typical assay contained 50 mM Hepes- Na^+ , pH 7.5, 100 mM NaCl, 10 μM CoCl_2 , 1 mM o-Dianisidine or 50 μM Amplex Red, 0.5 units of HRP (Sigma), 0.035 unit of AAO (Sigma), 1 nM hMetAP2, and varying amounts of peptide substrates. Assays were initiated by the addition of hMetAP2 enzyme, and the rates were corrected for the background rate determined in the absence of hMetAP2.



Kinetic Data Analysis:

Data were fitted to the appropriate rate equations using Grafit computer software. Initial velocity data conforming to Michaelis-Menton kinetics were fitted to eq. 4. Inhibition patterns conforming to apparent competitive and non-competitive inhibition were fitted to eq. 5 and eq. 6, respectively.

$$v = VA/(K_a + A) \quad (4)$$

$$v = VA/[K_a(1 + I/K_{is}) + A] \quad (5)$$

$$v = VA/[K_a(1 + I/K_{is}) + A(1 + I/K_{ii})] \quad (6)$$

- 10 In eqs. 4 - 6, v is the initial velocity, V is the maximum velocity, K_a is the apparent Michaelis constant, I is the inhibitor concentration, and A is the concentration of variable substrates. The nomenclature used in the rate equations for inhibition constants is that of Cleland (1963), in which K_{is} and K_{ii} represent the apparent slope and intercept inhibition constants, respectively.

Cell growth inhibition assays:

- 20 The ability of MetAP2 inhibitors to inhibit cell growth was assessed by the standard XTT microtitre assay. XTT, a dye sensitive to the pH change of mitochondria in eukaryotic cells, is used to quantify the viability of cells in the presence of chemical compounds. Cells seeded at a given number undergo approximately two divisions on average in the 72 hours of incubation. In the absence of any compound, this population of cells is in exponential growth at the end of the incubation period; the mitochondrial activity of these cells is reflected in the spectrophotometric readout (A450). Viability of a similar cell

population in the presence of a given concentration of compound is assessed by comparing the A450 reading from the test well with that of the control well. Flat-bottomed 96-well plates are seeded with appropriate numbers of cells ($4-6 \times 10^3$ cells/well in a volume of 200 μ l) from trypsinized exponentially growing cultures. In the case of HUVECs, the wells are coated with matrigel prior to establishing the cultures. To "blank" wells is added growth medium only. Cells are incubated overnight to permit attachment. Next day, medium from wells that contain cells is replaced with 180 μ l of fresh medium. Appropriate dilutions of test compounds are added to the wells, final DMSO concentration in all wells being 0.2 %. Cells plus compound are incubated for an additional 72 hr at 37°C under the normal growth conditions of the cell line used. Cells are then assayed for viability using standard XTT/PMS (prepared immediately before use: 8 mg XTT (Sigma X-4251) per plate is dissolved in 100 μ l DMSO. 3.9 ml H₂O is added to dissolve XTT and 20 μ l of PMS stock solution (30 mg/ml) is added from frozen aliquoted stock solution (10 mg of PMS (phenazine methosulfate, Sigma P-9625) in 3.3 ml PBS without cations. These stocks are frozen at -20°C until use). 50 μ l of XTT/PMS solution is added to each well and plates incubated for 90 minutes (time required may vary according to cell line, etc.) at 37°C until A₄₅₀ is >1.0. Absorbance at 450 nm is determined using a 96-well UV plate reader. Percent viability of cells in each well is calculated from these data (having been corrected for background absorbance). IC₅₀ is that concentration of compound that reduces cell viability to 50% control (untreated) viability.

The compounds of this invention show MetAP2 inhibitor activity having IC₅₀ values in the range of 0.0001 to 100 μ M. The full structure/activity relationship has not yet been established for the compounds of this invention. However, given the disclosure herein, one of ordinary skill in the art can utilize the present assays in order to determine which compounds of this invention are inhibitors of MetAP2 and which bind thereto with an IC₅₀ value in the range of 0.0001 to 100 μ M.

All publications, including, but not limited to, patents and patent applications cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.


The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments

specifically disclosed herein are within the scope of the following claims. Without further elaboration it is believed that one skilled in the art can, given the preceding description, utilize the present invention to its fullest extent. Therefore any examples are to be construed as merely illustrative and not a limitation on the scope of the

5 present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is claimed is:

1. A method of inhibiting MetAP2 in mammals, comprising administering to a mammal an effective amount of a compound of formula (IA) or a pharmaceutically acceptable salt or solvate thereof:



Formula (IA)

wherein,

X is S or O;

R¹ is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl;

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent.

2. The method of claim 1, wherein the compound of formula (IA) is selected from:

3-anilino-5-benzylthio-1,2,4-triazole;
 3-anilino-5-methylthio-1,2,4-triazole;
 3-anilino-5-(4-chloro-benzylthio)-1,2,4-triazole;
 3-anilino-5-allylthio-1,2,4-triazole;
 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole;
 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole;
 3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole;
 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-anilino-5-(propyl acetylthio)-1,2,4-triazole;
3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole;
3-anilino-5-(2-phenethylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(1*H*-benzoimidazol-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole;
3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole;
4-(5-anilino-2 *H*-[1,2,4]triazol-3-yl thio)-benzoic acid;
3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;

3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;

5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;

3-(2-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;

3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;

3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;

3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
[5-(benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3-methoxybenzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-oxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;

3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
: 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-methyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-butyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*i*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-allyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;

3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
 3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
 or a pharmaceutically acceptable salt or solvate thereof.

3. The method of claim 1, wherein the compound of formula (IA) is selected from:

3-anilino-5-benzylthio-1,2,4-triazole;
 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
 3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
 3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
 3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
 3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
 3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
 3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
 5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
 3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;

3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole
or a pharmaceutically acceptable salt or solvate thereof.

4. The method of claim 1, wherein the compound of formula (IA) is selected from:


3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

5. The method of claim 1, wherein the compound of formula (IA) is selected from:

3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

6. A method for treating a disease mediated by MetAP2 in mammals, comprising administering to a mammal in need of such treatment, an effective amount of a compound of formula (IA) or a pharmaceutically acceptable salt or solvate thereof:



Formula (IA)

wherein,

X is S or O;

R¹ is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent.

7. The method of claim 6, wherein the compound of formula (IA) is selected from:

- 3-anilino-5-benzylthio-1,2,4-triazole;
- 3-anilino-5-methylthio-1,2,4-triazole;
- 3-anilino-5-(4-chloro-benzylthio)-1,2,4-triazole;
- 3-anilino-5-allylthio-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole;
- 3-anilino-5-(α-methylbenzylthio)-1,2,4-triazole;
- 3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-anilino-5-(propyl acetylthio)-1,2,4-triazole;
3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole;
3-anilino-5-(2-phenethylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(1*H*-benzoimidazol-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole;
3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole;
4-(5-anilino-2 *H*-[1,2,4]triazol-3-yl thio)-benzoic acid;
3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
[5-(benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3-methoxybenzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-oxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;

3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
: 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-methyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-butyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*i*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-allyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;

3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
or a pharmaceutically acceptable salt or solvate thereof.

8. The method of claim 6, wherein the compound of formula (IA) is selected from:

3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;

3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

9. The method of claim 6, wherein the compound of formula (IA) is selected from:


3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

10. The method of claim 6, wherein the compound of formula (IA) is selected from:

3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

11. A method of treating conditions mediated by angiogenesis, selected from cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity, comprising administering a compound of formula (IA):



Formula (IA)

wherein:

X is S or O;

R¹ is optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent.

12. The method of claim 11, wherein the compound of formula (IA) is selected from:

- 3-anilino-5-benzylthio-1,2,4-triazole;
- 3-anilino-5-methylthio-1,2,4-triazole;
- 3-anilino-5-(4-chloro-benzylthio)-1,2,4-triazole;
- 3-anilino-5-allylthio-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-butylthio)-1,2,4-triazole;
- 3-anilino-5-(2-methyl-2-pentenylthio)-1,2,4-triazole;

3-anilino-5-(α -methylbenzylthio)-1,2,4-triazole;
3-anilino-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-anilino-5-(propyl acetylthio)-1,2,4-triazole;
3-anilino-5-(3,3-dimethoxy-propylthio)-1,2,4-triazole;
3-anilino-5-(2-phenethylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(3-phenyl-[1,2,4]oxadiazol-5-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(1*H*-benzoimidazol-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-(4-chlorophenyl)-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(4-*i*-propyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-anilino-5-(quinolin-8-ylthio)-1,2,4-triazole;
3-anilino-5-(4-acetamido-benzylthio)-1,2,4-triazole;
4-(5-anilino-2 *H*-[1,2,4]triazol-3-yl thio)-benzoic acid;
3-anilino-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-trifluoromethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-anilino-5-(3,5-dimethyl-benzylthio)-1,2,4-triazole;
3-anilino-5-(4-cyano-benzylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(4-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;

3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(cyclohexylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-2-butenylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(5-methyl-isoxazol-3-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(2-methyl-thiazol-4-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

4-(5-(pyridin-2-ylmethylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;

3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;

3-(3,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-benzylthio-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;

3-(2-phenyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(pyridin-4-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-phenyl-anilino)-5-(2-methyl-thiazol-4-ylmethylthio)-1,2,4-triazole;
[5-(benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3-methoxybenzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(cyclohexylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-2-butenylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-fluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(5-methyl-isoxazol-3-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(pyridin-2-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methyl-thiazol-4-ylmethylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1H-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;

3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;

3-(2-isopropyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-oxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;

3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;

3-(2-methyl-4-methoxy-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-methoxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-methylenedioxy-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(5-methyl-isoxazol-3-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(pyridin-2-ylmethylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
: 3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(cyclohexylmethylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-2-butenylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-methyl-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-chloro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-methyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*n*-butyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-*i*-propyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-allyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-benzyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;

3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
or a pharmaceutically acceptable salt or solvate thereof.

13. The method of claim 11, wherein the compound of formula (IA) is selected from:

3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole;
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;

3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;

3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole
or a pharmaceutically acceptable salt or solvate thereof.

14. The method of claim 11, wherein the compound of formula (IA) is selected from:


3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;

3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

15. The method of claim 11, wherein the compound of formula (IA) is selected from:

3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(2-isopropylphenylamino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-(2-isopropyl amino)-4H-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.

16. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof:



Formula (I)

wherein,

X is S or O;

R¹ is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl;

R² is optionally substituted C₂₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, provided that when R² is optionally substituted Het-C₀₋₆alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiazole or benzopyrazole, then the optional substituent is not -(CH₂)₂NR⁴R⁵; and

R³ is H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, or C₃₋₇cycloalkyl-C₀₋₆alkyl, C₀₋₆alkyl-C(O)X'AB, C₀₋₆alkyl-S(O)₂X'AB, C₀₋₆alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁₋₆alkyl, C₃₋₆alkenyl, C₃₋₆alkynyl, optionally substituted Ar-C₀₋₆alkyl, optionally substituted Het-C₀₋₆alkyl, C₃₋₇cycloalkyl-C₀₋₆alkyl, or A or B are independently absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

17. A pharmaceutical composition comprising a compound as claimed in claim 6 and a pharmaceutically acceptable carrier.

18. The compound of claim 16 which is selected from:
3-(4-chloro-anilino)-5-benzylthio-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;

3-(2-ethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(3-methyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-benzylthio-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methyl)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(*p*-methoxy)-anilino-5-benzylthio-1,2,4-triazole;
3-methylacetate-3-(2,6-dimethyl)-anilino-5-benzylthio-1,2,4-triazole;
3-anilino-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
5-(5-phenylamino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*p*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(2-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-*o*-tolyl amino-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(2-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;

3-(4-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
5-(5-(2-methoxyphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester
3-(2-methoxy-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(3-methyl-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(3-methylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-(4-*n*-butyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
3-(4-fluoro-anilino)-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
5-(5-(4-fluorophenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
3-anilino-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;

3-(2-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
4-(5-(3-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
methyl ester;
4-(5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
methyl ester;
4-(5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid
methyl ester;
3-(3,4-dimethoxy-anilino)-5-(3-methoxy-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3,4-dimethoxy-anilino)-5-(2-methoxy-benzylthio)-1,2,4-triazole;
[5-(2-fluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(3,4-difluoro-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(2-methoxy-benzylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
[5-(thiophen-2-ylthio)-1*H*-[1,2,4]triazol-3-yl]-pyridin-3-yl-amine;
3-(2-ethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-ethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(4-*n*-butyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,4-dimethoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;

3-(2-methyl-4-methoxy-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2-methyl-4-methoxy-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(4-fluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(3,4-difluoro-benzylthio)-1,2,4-triazole;
3-(2,6-dimethyl-anilino)-5-(2-fluoro-benzylthio)-1,2,4-triazole;
3-(4-fluoro-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole; and
3-(4-fluoro-anilino)-5-(furan-3-ylthio)-1,2,4-triazole
or a pharmaceutically acceptable salt or solvate thereof.

19. The compound of claim 16 which is selected from:

3-methylacetate-3-anilino-5-benzylthio-1,2,4-triazole;
4-(5-benzylthio-1*H*-[1,2,4]triazol-3-ylamino)-benzoic acid methyl ester;
3-anilino-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-3-ylthio)-1,2,4-triazole;
3-anilino-5-(furan-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-anilino-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-anilino-5-(5-bromo-thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
3-(4-methyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(4-chloro-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(4-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(2-methoxy-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
3-(3-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole; and
3-(3-methyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole,
or a pharmaceutically acceptable salt or solvate thereof.


20. The compound of claim 16 which is selected from:

3-(2-isopropyl-anilino)-5-(thiophen-2-ylthio)-1,2,4-triazole;
3-(2-isopropyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;


3-(2-isopropyl-anilino)-5-(3-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(5-methyl-thiophen-2-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole;
 5-(5-(2-isopropylphenylamino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carboxylic acid ethyl ester;
 5-(5-(2-isopropyl amino)-4*H*-[1,2,4]triazol-3-ylsulfanylmethyl)-furan-2-carbaldehyde;
 3-(2-isopropyl-anilino)-5-(thiophen-3-ylthio)-1,2,4-triazole;
 3-(2-isopropyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(furan-2-ylthio)-1,2,4-triazole;
 3-(4-methyl-anilino)-5-(furan-3-ylthio)-1,2,4-triazole;
 3-(2-methoxy-anilino)-5-(3-chloro-thiophen-2-ylthio)-1,2,4-triazole; and
 3-(2-methoxy-anilino)-5-(5-chloro-thiophen-2-ylthio)-1,2,4-triazole,
 or a pharmaceutically acceptable salt or solvate thereof.

21. A method of making a compound of formula (I), wherein X is sulfur, comprising the steps of:

a) combining an isothiocyanate of formula (II):




with thiourea to provide a compound of formula (III):




Formula (III)

b) treating the compound of formula (III) with an alkyl halide under basic conditions provide a compound of formula (IV):




Formula (IV)

c) treating the compound of formula (IV) with hydrazine to provide a compound of formula (V):




Formula (V)

d) treating the compound of formula (V) with an alkyl halide under basic conditions to provide a compound of formula (VI):




Formula (VI)

e) protecting the compound of formula (VI) to provide a compound of formula (VII):




Formula (VII)

f) alkylating the compound of formula (VII) with an alkyl halide to provide a compound of formula (VIII):



Formula (VIII)

g) deprotecting the compound of formula (VIII) to provide the compound of formula (I):



Formula (I)

wherein,

X is S;

R¹ is optionally substituted C₂-6alkyl, C₃-6alkenyl, C₃-6alkynyl, optionally substituted Ar-C₀-6alkyl, optionally substituted Het-C₀-6alkyl, or C₃-7cycloalkyl-C₀-6alkyl;

R² is optionally substituted C₂-6alkyl, C₃-6alkenyl, C₃-6alkynyl, optionally substituted Ar-C₀-6alkyl, optionally substituted Het-C₀-6alkyl, C₃-7cycloalkyl-C₀-6alkyl, provided that when R² is optionally substituted Het-C₀alkyl, and Het is indole, benzofuran, benzothiophene, benzisoxazole, benzothiozole or benzopyrazole, then the optional substituent is not -(CH₂)₂NR⁴R⁵; and

R³ is H, optionally substituted C₁-6alkyl, C₃-6alkenyl, C₃-6alkynyl, optionally substituted Ar-C₀-6alkyl, optionally substituted Het-C₀-6alkyl, or C₃-7cycloalkyl-C₀-6alkyl, C₀-6alkyl-C(O)X'AB, C₀-6alkyl-S(O)₂X'AB, C₀-6alkyl-X'AB, wherein X' is O, S, C or N; A and B are independently H, optionally substituted C₁-6alkyl, C₃-6alkenyl, C₃-6alkynyl, optionally substituted Ar-C₀-6alkyl, optionally substituted Het-C₀-6alkyl, C₃-7cycloalkyl-C₀-6alkyl, or A or B are independently absent, provided that the compound is not 5-anilino-3-benzylthio-1,2,4-triazole, 3-(4-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methyl-anilino)-5-benzylthio-1,2,4-triazole, 3-(4-methoxy-anilino)-5-benzylthio-1,2,4-triazole, 3-(2-methoxy-anilino)-5-benzylthio-1,2,4-triazole, or 3-ethyl-3-anilino-5-benzylthio-1,2,4-triazole.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/24796 A1

(51) International Patent Classification⁷: **A61K 31/422**,
31/4196, 31/4439, C07D 261/02, 249/12, 249/14, 401/02,
413/02

(74) Agents: **STEIN-FERNANDEZ**, Nora et al.; SmithKline
Beecham Corporation, Corporate Intellectual Property,
UW2220, 709 Swedeland Road, P.O. Box 1539, King of
Prussia, PA 19406-0939 (US).

(21) International Application Number: PCT/US00/26951

(22) International Filing Date:
29 September 2000 (29.09.2000)

(81) Designated States (*national*): AE, AL, AU, BA, BB, BG,
BR, BZ, CA, CN, CZ, DZ, EE, GE, GH, GM, HR, HU, ID,
IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK,
MN, MX, MZ, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT,
TZ, UA, US, UZ, VN, YU, ZA.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/157,286 1 October 1999 (01.10.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM CORPORATION
[US/US]; One Franklin Plaza, Philadelphia, PA 19103
(US).

Published:

- *With international search report.*
- *Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MARINO, Joseph,
P., Jr.** [US/US]; 214 Rock Glen Road, Wynnwood,
PA 19096 (US). **THOMPSON, Scott, K.** [US/US]; 75
Guilford Circle, Phoenixville, PA 19460 (US). **VEBER,
Daniel, Frank** [US/US]; 290 Batleson Road, Ambler, PA
19002 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: COMPOUNDS AND METHODS

(57) Abstract: Compounds of this invention are non-peptide, reversible inhibitors of type 2 methionine aminopeptidase, useful in treating conditions mediated by angiogenesis, such as cancer, haemangioma, proliferative retinopathy, rheumatoid arthritis, atherosclerotic neovascularization, psoriasis, ocular neovascularization and obesity.

WO 01/24796 A1

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

"Compounds and Methods"

the specification of which (check one)

☐ as attached hereto.

☒ as filed on **29 September 2000** as Serial No. **PCT/US00/26951**

☐ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or Inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Filing Date	Priority Claimed
--------	---------	-------------	------------------

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

Application Number	Filing Date
60/157,286	01 October 1999

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT

International application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Serial No.	Filing Date	Status
------------	-------------	--------

I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 20462.

Address all correspondence and telephone calls to Nora Stein-Fernandez, SmithKline Beecham Corporation, Corporate Intellectual Property-U.S., UW2220, P.O. Box 1539, King of Prussia, Pennsylvania 19406-0939, whose telephone number is 610-270-5044.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00 Full Name of Inventor: JOSEPH P. MARINO, JR.

Inventor's Signature: 

Date: 3-21-02

Residence: 709 Swedeland Road, King of Prussia, Pennsylvania 19406, United States of America

Citizenship: United States of America

Post Office Address: GlaxoSmithKline
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

2-00 Full Name of Inventor: SCOTT K. THOMPSON

Inventor's Signature: *Scott K. Thompson*

Date: 3-20-02

Residence: 709 Swedeland Road, King of Prussia, Pennsylvania 19406, United States of America

Citizenship: United States of America

Post Office Address: GlaxoSmithKline
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

3-00 Full Name of Inventor: DANIEL FRANK VEBER

Inventor's Signature: *Daniel Frank Veber*

Date: 3-22-02

Residence: 290 Batleson Road, Ambler, Pennsylvania 19002, United States of America

Citizenship: United States of America

Post Office Address: GlaxoSmithKline
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939